

THE CIRCLE

www.sesgeorgiaworks.com

Contribution to Society: Our People at Work

Mistaken Identity: Snakes



CASES OF MISTAKEN IDENTITY

A comparison of commonly confused Northeastern venomous and non-venomous snakes

VENOMOUS

NON-VENOMOUS





Crotatus horridus

- Recognized by a tall rattle, a shovel head, facial pits (see picture), and vertical pupils. Adult length: 36-54 inches
- Typically a yellowish color with brown or black bands down its back. Found in deciduous forests on rugged
- Hibernates in rocky crevices and dens.
- Endangered in several I NJ, VT, MA, and NH.





Eastern Hognose Heterodon platirhinos

Distinguished by an upturned snout (see

- picture), lack of facial pits, and round Adult length: 18-45 inches
- Variable coloration; can be spotted yellow or all black.
- Found in woodlands with dry sandy areas. When threatened the hognose will flatten its
- neck and bring its head up off the ground.
- It may also play dead.



Copperhead

- Recognized by a copper/brown/red shovel head, facial pits (see picture), and vertical
- Adult length: 24-36 inches Juveniles have tails with yellow tips.
- Found in forests, rocky outcrops, and farmland/fields.
- Endangered in MA







head, tapered tail, a lack of facial pits, and rounded pupils.

Adult length: 24-36 inches Typically has bands of red/brown/black

Distinguished by a Y or V on the top of its

Eastern Milk Snake

- down its body. Its underbelly is checkered white and black.
- Found in fields, forests, or rocky outcrops.
- Has a broad range across the Northeast.



- Recognized by black "hourglass" bands on its back, a flattened head, facial pits (see picture), and vertical pupils.
- Adult length: 30-48 inches When threatened the cottonmouth wil
- expose a white mouth Found near freshwater.
- Range does not extend further north than Virginia.







Northern Water Snake

Nerodia sipedon

- Distinguished by dark bands down its body, a narrow head, a slender body, a lack of facial pits, and round pupils
- Adult length: 24-42 inches Coloration can be brown, redish, gray, or
- Found near freshwater.
- Has a broad range across the Northeast.

- Venomous snakes are very rare in the Northeast Snakes are shy creatures and will typically not attack unless disturbed.
- Venom is meant for prey, not protection.
- Non-venomous snakes will also vibrate their tails when agitated.
- All Northeastern species of venomous snakes have heat sensing facial pits and vertical pupils.



- Snakes are an essential part of ecosystems because they act as both
- predator and prey. Their absence can alter the balance of a food web. Snakes control populations of smaller animals. This directly impacts humans because smaller mammals can spread disease, for example the

Poster Design by Erin Capra

